



VOLUME 6 - ANNEXES SECTION 6.28 – Espèces aviaires Rapport final de l'étude de faisabilité Phase I





Référence du consultant: LGA-1-GN-F-FRN-RT-0006_01_Section6.28



Stantec DESFOR SYSTIA

Liste des espèces aviaires rapportées sur le territoire Eeyou Istchee Baie-James

Source : Commission régionale sur les ressources naturelles et le territoire de la Baie-James, 2010.

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|-----------------------------|--------------------|--------------------------------|---------------------------------------------------------------|
| Oie des neiges | | Chen caerulescens | Present during migration |
| Grande oie des neiges | | Chen caerulescens atlantica | Present during migration |
| Petite oie des neiges | | Chen cearulescens caerulescens | Present during migration |
| Bernache cravant | | Branta bernicla | Present during migration |
| Bernache nonnette | | Branta leucopsis | (Rare/Accidental) |
| Bernache du Canada | | Branta canadensis | Present during summer |
| Bernache de Hutchins | | Branta hutshinsii | Present during migration |
| Cygne siffleur | | Cygnus columbianus | Present during migration |
| Canard branchu | | AiX sponsa | Present during summer |
| Canard chipeau | | Anas strepera | Present during migration (Rare/Accidental) |
| Canard d'Amérique | | Anas americana | Present during summer |
| Canard noir | | Anas rubripes | Present during summer |
| Canard colvert | | Anas platyrhynchos | Present during summer |
| Hybride canard colvert-noir | | - | Present during summer |
| Sarcelle à ailes bleues | | Anas discors | Present during summer |
| Canard souchet | | Anas clypeata | Present during summer |
| Canard pilet | | Anas acuta | Present during summer |
| Sarcelle d'hiver | | Anas crecca | Present during summer |
| Fuligule à dos blanc | | Aythya valisineria | Present south of territory during migration (Rare/Accidental) |
| Fuligule à tête rouge | | Aythya americana | Present south of territory during migration (Rare/Accidental) |
| Fuligule à collier | | Aythya collaris | Present during summer |
| Fuligule milouinan | | Aythya marila | Present during summer and migration |
| Petit Fuligule | | Aythya affinis | Present during summer |
| Eider à duvet | | Somateria mollissima | Present during summer |
| Macreuse à front blanc | | Melanitta perspicillata | Present during summer |
| Macreuse brune | | Melanitta fusca | Present during summer |
| Macreuse noire | | Melanitta nigra | Present during migration |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|-------------------------|--------------------|---------------------------|-------------------------------------------------------|
| Harelde kakawi | | Clangula hyemalis | Present during summer and migration |
| Arlequin plongeur | | Histrionicus histrionicus | Present during summer |
| Petit Garrot | | Bucephala albeola | Present during summer |
| Garrot à œil d'or | | Bucephala clangula | Present during summer |
| Garrot d'Islande | | Bucephala islandica | Present during summer |
| Harle couronné | | Lophodytes cucullatus | Present during summer |
| Grand Harle | | Mergus merganser | Present during summer |
| Harle huppé | | Mergus serrator | Present during summer |
| Érismature rousse | | OXyura jamaicensis | Present during summer and migration (Rare/Accidental) |
| Gélinotte huppée | | Bonasa umbellus | Present during summer |
| Tétras du Canada | | Falcipennis canadensis | Present during summer |
| Lagopède des saules | | Lagopus lagopus | Present during winter |
| Lagopède alpin | | Lagopus muta | Present during winter |
| Tétras à queue fine | | Tympanuchus phasianellus | Resident |
| Plongeon catmarin | | Gavia stellata | Present during summer |
| Plongeon du Pacifique | | Gavia pacifica | Present during summer |
| Plongeon huard | | Gavia immer | Present during summer |
| Grèbe à bec bigarré | | Podilymbus podiceps | Present during summer (Rare/Accidental) |
| Grèbe esclavon | | Podiceps auritus | Present during summer (Rare/Accidental) |
| Grèbe jougris | | Podiceps grisegena | Present during summer (Rare/Accidental) |
| Cormoran à aigrettes | | PhalacrocoraX auritus | Present during summer |
| Butor d'Amérique | | Botaurus lentiginosus | Present during summer |
| Grand Héron | | Ardea herodias | Present during summer |
| Bihoreau gris | | NycticoraX nycticoraX | Present during summer (Rare/Accidental) |
| Urubu à tête rouge | | Cathartes aura | Present during summer (Rare/Accidental) |
| Urubu noir | | Coragyps atratus | Present during summer (Rare/Accidental) |
| Balbuzard pécheur | | Pandion haliaetus | Present during summer |
| Pygargue à tête blanche | | Haliaeetus leucocephalus | Present during summer |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|---------------------------|--------------------|-----------------------------|--------------------------------------------|
| Busard Saint-Martin | | Circus cyaneus | Present during summer |
| Épervier brun | | Accipiter striatus | Present during summer |
| Autour des palombes | | Accipiter gentilis | Resident |
| Petite Buse | | Buteo platypterus | Present during summer (Rare/Accidental) |
| Buse à queue rousse | | Buteo jamaicensis | Present during summer |
| Buse pattue | | Buteo lagopus | Present during summer |
| Aigle royal | | Aquila chrysaetos | Present during summer |
| Crécerelle d'Amérique | | Falco sparverius | Present during summer |
| Faucon émerillon | | Falco columbarius | Present during summer |
| Faucon gerfaut | | Falco rusticolus | Present during winter |
| Faucon pèlerin (anatum) | | Falco peregrinus anatum | Present during migration |
| Faucon pèlerin (tundrius) | | Falco peregrinus tundrius | Present during migration |
| Râle jaune | | Coturnicops noveboracensis | Present during summer |
| Marouette de Caroline | | Porzana carolina | Present during summer |
| Gallinule poule-d'eau | | Gallinula chloropus | Present during summer (Rare/Accidental) |
| Foulque d'Amérique | | Fulica americana | Present during summer |
| Grue du Canada | | Grus canadensis | Present during summer |
| Pluvier argenté | | Pluvialis squatarola | Present during migration |
| Pluvier bronzé | | Pluvialis dominica | Present during migration |
| Pluvier semipalmé | | Charadrius semipalmatus | Present during summer and migration |
| Pluvier kildir | | Charadrius vociferus | Present during summer |
| Chevalier grivelé | | Actitis macularius | Present during summer |
| Chevalier solitaire | | Tringa solitaria | Present during summer |
| Chevalier semipalmé | | Catoptrophorus semipalmatus | (Rare/Accidental) |
| Grand Chevalier | | Tringa melanoleuca | Present during summer and migration |
| Petit Chevalier | | Tringa flavipes | Present during summer and migration |
| Maubèche des champs | | Bartramia longicauda | Present during summer (Rare/Accidental) |
| Courlis esquimau | | Numenius borealis | Present during migration (Rare/Accidental) |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|------------------------------|--------------------|------------------------------|-------------------------------------------------------|
| Courlis corlieu | | Numenius phaeopus | Present during migration |
| Barge hudsonienne | | Limosa haemastica | Present during migration |
| Barge marbrée | | Limosa fedoa | Present during summer |
| Tournepierre à collier | | Arenaria interpres | Present during migration |
| Bécasseau maubèche | | Calidris canutus | Present during migration |
| Bécasseau sanderling | | Calidris alba | Present during migration |
| Bécasseau semipalmé | | Calidris pusilla | Present during summer and migration |
| Bécasseau minuscule | | Calidris minutilla | Present during summer and migration |
| Bécasseau à croupion blanc | | Calidris fuscicollis | Present during migration |
| Bécasseau de Baird | | Calidris bairdii | Present during migration |
| Bécasseau à poitrine cendrée | | Calidris melanotos | Present during migration |
| Bécasseau violet | | Calidris maritima | Present during migration (Rare/Accidental) |
| Bécasseau variable | | Calidris alpina | Present during migration |
| Bécasseau d'Alaska | | Calidris mauri | Present during migration (Rare/Accidental) |
| Bécasseau à échasse | | Calidris himantopus | Present during migration |
| Bécasseau roussâtre | | Tryngites subruficollis | Present during migration |
| Bécassin rouX | | Limnodromus griseus | Present during summer and migration |
| Bécassin à long bec | | Limnodromus scolopaceus | Present during migration |
| Bécasse d'Amérique | | ScolopaX minor | Present during summer |
| Bécassine de Wilson | | Gallinago delicata | Present during summer |
| Phalarope de Wilson | | Phalaropus tricolor | Present during summer and migration (Rare/Accidental) |
| Phalarope à bec étroit | | Phalaropus lobatus | Present during summer and migration |
| Phalarope à bec large | | Phalaropus fulicarius | (Rare/Accidental) |
| Mouette de Bonaparte | | Chroicocephalus philadelphia | Present during summer |
| Mouette pygmée | | Larus minutus | Present during summer |
| Mouette de Franklin | | Leucophaeus pipiXcan | Present during summer (Rare/Accidental) |
| Mouette tridactyle | | Rissa tridactyla | (Rare/Accidental) |
| Goéland à bec cerclé | | Larus delawarensis | Present during summer |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|---------------------------|--------------------|----------------------|-----------------------------------------|
| Goéland argenté | | Larus argentatus | Present during summer |
| Goéland arctique | | Larus glaucoides | Present during migration |
| Goéland bourgmestre | | Larus hyperboreus | Present during migration |
| Goéland marin | | Larus marinus | |
| Guifette noire | | Chlidonias niger | Present during summer (Rare/Accidental) |
| Sterne pierregarin | | Sterna hirundo | Present during summer |
| Sterne arctique | | Sterna paradisaea | Present during summer and migration |
| Sterne caspienne | | Sterna caspia | Present during summer (rare au Québec) |
| Guillemot à miroir | | Cepphus grylle | Resident |
| Pigeon biset | | Columba livia | Resident (low abundance) |
| Tourterelle triste | | Zenaida macroura | Present during summer |
| Grand-duc d'Amérique | | Bubo virginianus | Resident |
| Harfang des neiges | | Bubo scandiacus | Present during winter |
| Chouette épervière | | Surnia ulula | Resident |
| Chevêche des terriers | | Athene cunicularia | Present during summer (Rare/Accidental) |
| Chouette rayée | | StriX varia | Resident |
| Chouette lapone | | StriX nebulosa | Resident |
| Hibou moyen-duc | | Asio otus | Present during summer (Rare/Accidental) |
| Hibou des marais | | Asio flammeus | Present during summer |
| Nyctale de Tengmalm | | Aegolius funereus | Resident |
| Petite Nyctale | | Aegolius acadicus | Present during summer (Rare/Accidental) |
| Engoulevent d'Amérique | | Chordeiles minor | Present during summer |
| Martinet ramoneur | | Chaetura pelagica | Present during summer (Rare/Accidental) |
| Colibri à gorge rubis | | Archilochus colubris | Present during summer (Rare/Accidental) |
| Martin-pêcheur d'Amérique | | Megaceryle alcyon | Present during summer |
| Pic maculé | | Sphyrapicus varius | Present during summer |
| Pic mineur | | Picoides pubescens | Resident |
| Pic chevelu | | Picoides villosus | Resident |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|----------------------------|--------------------|------------------------|---------------------------------------------------------------------------------|
| Pic à dos rayé | | Picoides dorsalis | Resident |
| Pic à dos noir | | Picoides arcticus | Resident |
| Pic flamboyant | | Colaptes auratus | Present during summer |
| Grand pic | | Dryocopus pileatus | Resident (Rare/Accidental) |
| Moucherolle à côtés olive | | Contopus cooperi | Present during summer |
| Moucherolle à ventre jaune | | EmpidonaX flaviventris | Present during summer |
| Moucherolle des aulnes | | EmpidonaX alnorum | Present during summer |
| Moucherolle tchébec | | EmpidonaX minimus | Present during summer |
| Moucherolle phébi | | Sayornis phoebe | (Rare/Accidental) |
| Tyran tritri | | Tyrannus tyrannus | Present during summer |
| Tyran à longue queue | | Tyrannus forficatus | Present during summer (Rare/Accidental) |
| Pie-grièche grise | | Lanius eXcubitor | Present during summer and migration |
| Viréo à tête bleue | | Vireo solitarius | Present during summer |
| Viréo de Philadelphie | | Vireo philadelphicus | Present during summer |
| Viréo auX yeuX rouges | | Vireo olivaceus | Present during summer |
| Mésangeai du Canada | | Perisoreus canadensis | Resident |
| Geai bleu | | Cyanocitta cristata | Present during summer (Rare/Accidental) |
| Pie d'Amérique | | Pica hudsonia | (Rare/Accidental) |
| Corneille d'Amérique | | Corvus brachyrhynchos | Present during summer |
| Grand Corbeau | | Corvus coraX | Resident |
| Alouette hausse-col | | Eremophila alpestris | Present during summer and migration |
| Hirondelle bicolore | | Tachycineta bicolor | Present during summer |
| Hirondelle de rivage | | Riparia riparia | Present during summer |
| Hirondelle rustique | | Hirundo rustica | Present during summer |
| Mésange à tête noire | | Pœcile atricapillus | Resident |
| Mésange à tête brune | | Pœcile hudsonicus | Resident |
| Sittelle à poitrine rousse | | Sitta canadensis | Present during summer and during summer south of 50th parallel approximatively. |
| Grimpereau brun | | Certhia americana | Present during summer |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|---------------------------|--------------------|-------------------------|---------------------------------------------------------------------------------------|
| Troglodyte mignon | | Troglodytes troglodytes | Present during summer |
| Roitelet à couronne dorée | | Regulus satrapa | Present during summer |
| Roitelet à couronne rubis | | Regulus calendula | Present during summer |
| Merlebleu de l'Est | | Sialia sialis | Present during summer (Rare/Accidental) |
| Grive à joues grises | | Catharus minimus | Present during summer and migration |
| Grive à dos olive | | Catharus ustulatus | Present during summer |
| Grive solitaire | | Catharus guttatus | Present during summer |
| Grive fauve | | Catharus fuscescens | Present during summer |
| Merle d'Amérique | | Turdus migratorius | Present during summer |
| Moqueur polyglotte | | Mimus polyglottos | Present during summer (Rare/Accidental) |
| Étourneau sansonnet | | Sturnus vulgaris | Present during summer |
| Pipit d'Amérique | | Anthus rubescens | Present during summer and migration |
| Jaseur boréal | | Bombycilla garrulus | Present during winter south of 52th parallel and during summer north of 52th parallel |
| Jaseur d'Amérique | | Bombycilla cedrorum | Present during summer |
| Paruline obscure | | Vermivora peregrina | Present during summer |
| Paruline verdâtre | | Vermivora celata | Present during summer |
| Paruline à joues grises | | Vermivora ruficapilla | Present during summer |
| Paruline jaune | | Dendroica petechia | Present during summer |
| Paruline à flancs marron | | Dendroica pensylvanica | Present during summer |
| Paruline à tête cendrée | | Dendroica magnolia | Present during summer |
| Paruline tigrée | | Dendroica tigrina | Present during summer |
| Paruline à croupion jaune | | Dendroica coronata | Present during summer |
| Paruline à gorge noire | | Dendroica virens | Present during summer |
| Paruline à gorge orangée | | Dendroica fusca | Present during summer (Rare/Accidental) |
| Paruline à joues grises | | Vermivora ruficapilla | Present during summer |
| Paruline jaune | | Dendroica petechia | Present during summer |
| Paruline à flancs marron | | Dendroica pensylvanica | Present during summer |
| Paruline à tête cendrée | | Dendroica magnolia | Present during summer |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|----------------------------|--------------------|---------------------------|--------------------------------------------|
| Paruline tigrée | | Dendroica tigrina | Present during summer |
| Paruline à croupion jaune | | Dendroica coronata | Present during summer |
| Paruline à gorge noire | | Dendroica virens | Present during summer |
| Paruline à couronne rousse | | Dendroica palmarum | Present during summer |
| Paruline à poitrine baie | | Dendroica castanea | Present during summer |
| Paruline rayée | | Dendroica striata | Present during summer |
| Paruline noir et blanc | | Mniotilta varia | Present during summer |
| Paruline flamboyante | | Setophaga ruticilla | Present during summer |
| Paruline couronnée | | Seiurus aurocapilla | Present during summer |
| Paruline des ruisseauX | | Seiurus noveboracensis | Present during summer |
| Paruline à gorge grise | | Oporornis agilis | Present during summer |
| Paruline triste | | Oporornis philadelphia | Present during summer |
| Paruline masquée | | Geothlypis trichas | Present during summer |
| Paruline à calotte noire | | Wilsonia pusilla | Present during summer |
| Paruline du Canada | | Wilsonia canadensis | Present during summer |
| Bruant hudsonien | | Spizella arborea | Present during summer and migration |
| Bruant familier | | Spizella passerina | Present during summer |
| Bruant vespéral | | Pooecetes gramineus | Present during summer (Rare/Accidental) |
| Bruant des prés | | Passerculus sandwichensis | Present during summer |
| Bruant de Le Conte | | Ammodramus leconteii | Present during summer (Rare/Accidental) |
| Bruant de Nelson | | Ammodramus nelsoni | Present during summer |
| Bruant fauve | | Passerella iliaca | Present during summer and migration |
| Bruant chanteur | | Melospiza melodia | Present during summer |
| Bruant de Lincoln | | Melospiza lincolnii | Present during summer |
| Bruant des marais | | Melospiza georgiana | Present during summer |
| Bruant à gorge blanche | | Zonotrichia albicollis | Present during summer |
| Bruant à face noire | | Zonotrichia querula | Present during migration (Rare/Accidental) |
| Bruant à couronne blanche | | Zonotrichia leucophrys | Present during summer and migration |

| Nom commun français | Nom commun anglais | Nom scientifique | Occurrence saisonnière |
|-----------------------|--------------------|-------------------------------|----------------------------------------------------------------------------------|
| Junco ardoisé | | Junco hyemalis | Present during summer |
| Bruant lapon | | Calcarius lapponicus | Present during summer and migration |
| Bruant de Smith | | Calcarius pictus | Present during migration (Rare/Accidental) |
| Bruant des neiges | | PlectrophenaX nivalis | Present during migration |
| Sturnelle des prés | | Steunella magna | Present during summer (Rare/Accidental) |
| Carouge à épaulettes | | Agelaius phoeniceus | Present during summer |
| Carouge à tête jaune | | Xanthocephalus Xanthocephalus | Present during summer |
| Quiscale rouilleuX | | Euphagus carolinus | Present during summer |
| Quiscale bronzé | | Quiscalus quiscula | Present during summer |
| Vacher à tête brune | | Molothrus ater | Present during summer |
| Durbec des sapins | | Pinicola enucleator | Resident south of 51th parallel and during summer north of 51th parallel |
| Roselin pourpré | | Carpodacus purpureus | Present during summer |
| Bec-croisé des sapins | | LoXia curvirostra | Resident (Rare/Accidental) |
| Bec-croisé bifascié | | LoXia leucoptera | Resident |
| Sizerin flammé | | Acanthis flammea | Present during winter south of 52th parallel and resident north of 52th parallel |
| Sizerin blanchâtre | | Acanthis hornemanni | Present during winter |
| Tarin des pins | | Spinus pinus | Resident south of 50th parallel and during summer north of 50th parallel |
| Chardonneret jaune | | Spinus tristis | Present during summer |
| Chardonneret élégant | | Carduelis carduelis | Present during summer (non native) |
| Gros-bec errant | | Coccothraustes vespertinus | Present during summer |
| Moineau domestique | | Passer domesticus | Resident (non native) |